

農学工学総合研究科博士後期課程課程3年の津田浩利氏（主指導教員：國武教授）の投稿論文が HortScience 49 巻 2 号（2014 年 2 月）に掲載され、表紙に採用されました。

Cover Photo — February 2014, 49 (2)

1/1 

Cover image



ON THE COVER

Tsuda et al. tested efficient *in vitro* methods for screening the genotypes with higher pH tolerance using multiple shoots of intersectional hybrids between *Vaccinium corymbosum* Spartan and *V. bracteatum*. The cover shows the effect of medium pH on root viability of two intersectional hybrids (JM1 and JM4) at Spartan. Roots of the *in vitro* plantlets were soaked in liquid media with different pH levels for 6 h. Then roots were stained with fluorescein diacetate/propidium iodide solution. For more information, see the paper beginning on p. 141.

HortScience

Volume 49, Number 2

CONTENTS

February 2014

COLLOQUIUM

- 112 Progress Toward Increasing Intake of Dietary Nutrients from Vegetables and Fruits: The Case for a Greater Role for the Horticultural Sciences
Philipp W. Simon
- 116 The Intersection of Plant Breeding, Human Health, and Nutritional Security: Lessons Learned and Future Perspectives
Bhimanagouda S. Patil, Kevin Crosby, David Byrne, and Kendal Hirachi
- 128 Assessing Nutritional Changes in a Vegetable Over Time: Issues and Considerations
Mark W. Furnham and Michael A. Gruak
- 133 The Future of Breeding Vegetables with Human Health Functionality: Realities, Challenges, and Opportunities
Irwin L. Goldman

REPORTS

- Breeding, Cultivars, Rootstocks, and Germplasm Resources**
- 138 Fruit Quality of Pear Psylla-resistant Parental Germplasm
Richard L. Bell
- 141 Efficient In Vitro Screening for Higher Soil pH Adaptability of Intersectional Hybrids in Blueberry
Hirotoshi Tsuda, Hisato Kunitake, Yo Aoki, Akiko Oyama, Takuya Tetsumura, Haruki Komatsu, and Katsunori Yoshioka
- 145 Identification, Nomenclature, Genome Sizes, and Ploidy Levels of *Liriope* and *Ophiopogon* Taxa
Jason D. Lattier, Thomas G. Ranney, Paul R. Fantz, and Tony Avent
- Crop Production**
- 152 Controlled-release Fertilizer during Cut Affects Growth and Tissue Nutrient Co of Rooted Cuttings of Annual Bedding
Christopher J. Currey and Roberto C

- 160 The Number of Emitters Alters Salt Distribution and Root Growth in Potted Gerbers
Raquel Valdés, Julián Miralles, Jesús Ochoa, and María del Mar García

Soil Management, Fertilization, and Irrigation

- 201 Hydroponic Production of Purslane as a Sodium-removing Vegetable in NaCl-rich Nutrient Solution
Yun Kong and Youbin Zheng
- 207 Statistical Model for Describing Macronutrient Impacts on Container Substrate pH Over Time
Jared Barnes, Paul Nelson, Brian E. Whipker, David A. Dickey, Dean Heisterberg, and Wei Shi
- 215 Interaction of Irrigation and Soil Management on Sweet Cherry Productivity and Fruit Quality at Different Crop Loads that Simulate Those Occurring by Environmental Extremes
Gerry H. Nielsen, Denise Nielsen, Frank Kappel, and T. Forge

Turf Management

- 221 Drought Resistance Strategies of Seashore Paspalum Cultivars at Different Mowing Heights
Mohamed A. Shabba, Mohamed S. Abbas, and Saad F. Alshammary

MISCELLANEOUS

- 230 Volatile Toluene and Xylene Removal Efficiency of Foliage Plants as Affected by Top to Root Zone Size
Kwang Jin Kim, Hyun Hwan Jung, Hyo Won Seo, Jung A. Lee, and Stanley J. Kays
- 235 Reported Death
- 235 Corrigendum

ON THE COVER

Tsuda et al. tested efficient in vitro methods for screening the genotypes with higher pH tolerance using multiple shoots of intersectional hybrids between *Faccinium corymbosum* Spartan and *F. bracteatum*. The cover shows the effect of medium pH on root viability of two intersectional hybrids (JM1 and JM4) and Spartan. Roots of the in vitro plantlets were soaked in liquid media with different pH levels for 6 h. Then roots were stained with fluorescein diacetate/propidium iodide solution. For more information, see the paper beginning on p. 141.



Image Analysis Systems for PI

Need More Than One Software Program?

Take Advantage of Our New Suites.



110

ON THE COVER

Tsuda et al. tested efficient in vitro methods for screening the genotypes with higher pH tolerance using multiple shoots of intersectional hybrids between *Faccinium corymbosum* Spartan and *F. bracteatum*. The cover shows the effect of medium pH on root viability of two intersectional hybrids (JM1 and JM4) and Spartan. Roots of the in vitro plantlets were soaked in liquid media with different pH levels for 6 h. Then roots were stained with fluorescein diacetate/propidium iodide solution. For more information, see the paper beginning on p. 141.

